

New Limits on Sterile Neutrino Mixing with Atmospheric Neutrinos

Alex Himmel

Duke University

for the

The Super Kamiokande Collaboration


Duke
UNIVERSITY



13th International Conference on Topics in
Astroparticle and Underground Physics


September 12th, 2013

What is a Sterile Neutrino?





What is a Sterile Neutrino?

3 neutrinos \rightarrow 2 mass splittings



What is a Sterile Neutrino?




3 neutrinos \rightarrow 2 mass splittings



3 mass splittings \rightarrow 4 neutrinos

What is a Sterile Neutrino?

3 neutrinos \rightarrow 2 mass splittings



3 mass splittings \rightarrow 4 neutrinos

One of which does not couple to the Z and so does not interact weakly, i.e. **sterile**

What is a Sterile Neutrino?



PMNS

$$U = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} & U_{e4} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} & U_{\mu 4} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} & U_{\tau 4} \\ U_{s1} & U_{s2} & U_{s3} & U_{s4} \end{pmatrix}$$

- One more neutrino adds 7 complex matrix elements
 - but not all independent.
- 1 Δm^2 , 3 “angles”, 2 phases – varying parameterizations
 - $|U_{e4}|^2, |U_{\mu 4}|^2, |U_{\tau 4}|^2$ or $\theta_{14}, \theta_{24}, \theta_{34}$
- One more neutrino adds **8 more parameters**

Evidence of Sterile Neutrinos

$$\Delta m^2 \approx 0.1 - 10 \text{ eV}^2$$




2-flavor
approx.

$$\sin^2 2\theta_{\mu e} = 4|U_{e4}|^2|U_{\mu 4}|^2$$

Both $|U_{e4}|^2$ and $|U_{\mu 4}|^2$ must be > 0
for non-zero probability

PRL110, 161801 (2013)

Evidence of Sterile Neutrinos



2-flavor
approx.

$$\sin^2 2\theta_{ee} = 4|U_{e4}|^2(1 - |U_{e4}|^2)$$

What can Super-K tell us?

- 11 years of atmospheric neutrino data.
 - Covering a wide range of L and E
- It is most useful because of **what it is not sensitive to:**

The size of the sterile mass splitting




Oscillations appear “fast”

The number of sterile neutrinos




3+1 and 3+N models look the same



What can Super-K tell us?

- Short-baseline-related: $|U_{\mu 4}|^2$
 - Driven by new Δm^2
 - Creates fast oscillations across a wide range of ν_μ samples
- Atmospheric/long-baseline: $|U_{\tau 4}|^2$
 - Accessible *only* at long distances
 - Oscillations into ν_s instead of ν_τ
 - Introduces a new matter effect




Super-K Sterile Model


- A fully generic sterile model **is difficult** computationally
 - Cannot calculate both active (ν_e) and sterile (NC) matter effects together
- So, we need to perform 2 different fits:

Fit for $ U_{\mu 4} ^2$	Fit for $ U_{\tau 4} ^2$
<ul style="list-style-type: none">– ν_e matter effects only– Most accurate $U_{\mu 4} ^2$ limit– No $U_{\tau 4} ^2$ limit	<ul style="list-style-type: none">– NC matter effects only– Required for $U_{\tau 4} ^2$– Over-constrains $U_{\mu 4} ^2$



SK Analysis: Zenith Angle

- Look for change in flavor content vs. L
- Bin by angle and separate μ and e
 - isolate oscillations
 - other samples control systematics
- The original SK analysis was simple: up/down, μ/e







Fully Contained
Sub-GeV Multi-GeV



Partially
Contained



Up-going μ



100's of MeV

Few GeV

1 TeV+

μ -like, e -like ($\nu_e/\bar{\nu}_e$), $\text{NC}\pi^0$ -like

Low energy ->

Poor $\cos\theta_z$ resolution

Long tracks – all μ -like

Uncontained ->

Poor E resolution

Oscillogram: Standard 3v


FC Sub-GeV

Multi-GeV

PC, Stop μ

Through-going μ

$$P(\nu_\mu \text{ to } \nu_\mu)$$



Oscillogram:

$|U_{\mu 4}|^2$ Fit


FC Sub-GeV

Multi-GeV


PC, Stop μ

Through-going μ

$P(\nu_\mu \text{ to } \nu_\mu)$



Fit for $|U_{\mu 4}|^2$



- Signature is extra disappearance in all μ samples
 - Correlated change at all energies, all $\cos\theta_z$
 - Sensitivity limited by μ/e flux uncertainty


Need to do 2 fits since we cannot calculate ν_e and NC matter effects simultaneously

Fit for $|U_{\mu 4}|^2$

$|U_{\mu 4}|^2 < 0.023$ at 90% C.L.

$|U_{\mu 4}|^2 < 0.034$ at 99% C.L.

As with similar experiments, no sterile-driven ν_μ disappearance



Need to do 2 fits since we cannot calculate ν_e and NC matter effects simultaneously

PRD86, 052009 (2012)

PRL52, 1384 (1984)


Fit for $|U_{\mu 4}|^2$

$|U_{\mu 4}|^2 < 0.023$ at 90% C.L.

$|U_{\mu 4}|^2 < 0.034$ at 99% C.L.

As with similar
experiments, no sterile-
driven ν_μ disappearance

Exclude much of the
MiniBooNE appearance
signal



Need to do 2 fits since we cannot calculate
 ν_e and NC matter effects simultaneously

PRD86, 052009 (2012)

ArXiv:1303.3011

Oscillogram:

$|U_{\tau 4}|^2$ Fit

FC Sub-GeV


Multi-GeV

PC, Stop μ


Through-going μ

$P(\nu_\mu \text{ to } \nu_\mu)$

Potentially
large changes
introduced by
sterile/NC
matter effects



Fit for $|U_{\tau 4}|^2$ (with $|U_{\mu 4}|^2$)



5-100 GeV

- Matter effects create shape distortion in PC/Up- μ zenith distribution
 - Less disappearance in most upward bins, still have extra disappearance in downward bins

Need to do 2 fits since we cannot calculate ν_e and NC matter effects simultaneously


Fit for $|U_{\tau 4}|^2$ (with $|U_{\mu 4}|^2$)

$|U_{\tau 4}|^2 < 0.28$ at 99% C.L.

Favors μ to τ oscillations
over μ to s

Lack of sterile matter effects
places a strong constraint

- Note, $|U_{\mu 4}|^2$ is over-constrained in this fit



All comparisons from:
ArXiv:1303.3011


Conclusions

- Atmospheric neutrinos provide a useful tool to study sterile oscillations
 - Wide range of L/E
 - Measurement independent of Δm^2 and $N_{\text{sterile}} > 1$
- No evidence of sterile neutrinos seen
 - No sterile-driven ν_μ disappearance, consistent with other short- and long-baseline measurements
 - μ to s oscillations strongly disfavored by the lack of sterile matter effects

3+N \approx 3+1 for Super K


$$P_{\mu\mu} = \boxed{(1 - |U_{\mu 4}|^2)^2} P_{\mu\mu}^0 + \boxed{\sum_{i \geq 4} |U_{\mu i}|^4}$$

- The first sterile term:
 - Controls extra disappearance
 - Is the same for any N_{sterile}
- The second sterile term:
 - Fills in the minima
 - Varies for N_{sterile}
- Our experiment is *much* more sensitive to **first term**
 - Beam experiments, focusing on the first oscillation dip, *are* sensitive to **the second term**.




No Matter in 3ν

- At right are 2 sensitivities from the $2+1$ fit
- The dashed is the normal fit, solid has sterile matter effects arbitrarily turned off
- $|U_{\mu 4}|^2$ limit is unaffected – it is independent of the sterile matter effects




When is Δm^2_{41} no longer “large”?

- When do the oscillations no longer appear fast?
 - This will be the worst at short L's and large E's, so let's focus on Up- μ with $\cos \theta_z > -0.1$
 - Loop through all these events and calculate the mean of $\sin^2(\Delta m^2 L/4E)$ for various Δm^2
- Doing this, the approximation is valid down to $\sim 0.8 \text{ eV}^2$



When is Δm^2_{41} no longer “large”?

- However, the limit on $|U_{\mu 4}|^2$ is driven by the low $|U_{\tau 4}|^2$ region.
 - In this region, the dominant samples are Sub-GeV muons
 - Almost no power comes from Up- μ
- For these samples, the “large” assumption is \sim always valid so $|U_{\mu 4}|^2$ limit really is a vertical line in Δm^2 to a good approximation



Our Matter Effect Model

- The μ to μ probability is fairly simple:

$$P_{\mu\mu} = (1 - d_\mu)^2 |\tilde{S}_{22}|^2 + d_\mu^2$$

- Get $|S_{22}|^2$ by diagonalizing the sum of the vacuum and matter Hamiltonians:

$$\begin{aligned} H^{(2)} &= H_{sm}^{(2)} + H_s^{(2)} : \\ &= \frac{\Delta m_{31}^2}{4E} \begin{pmatrix} -\cos 2\theta_{23} & \sin 2\theta_{23} \\ \sin 2\theta_{23} & \cos 2\theta_{23} \end{pmatrix} \pm \frac{G_F N_n}{\sqrt{2}} \begin{pmatrix} |\tilde{U}_{s2}|^2 & \tilde{U}_{s2}^* \tilde{U}_{s3} \\ \tilde{U}_{s2} \tilde{U}_{s3}^* & |\tilde{U}_{s3}|^2 \end{pmatrix} \end{aligned}$$

Our Matter Effect Model

- This gives us a \sim familiar matter-effect probability

$$|\tilde{S}_{22}|^2 = 1 - \sin^2(2\theta_m) \sin^2(f_m L)$$

$$f_m = \sqrt{A_{32}^2 + A_s^2 + 2A_{32}A_s (\cos(2\theta_{23}) \cos(2\theta_s) + \sin(2\theta_{23}) \sin(2\theta_s))}$$

$$E_{1,2}^m = \pm f_m$$

$$\sin 2\theta_m = \frac{A_{32} \sin(2\theta_{23}) + A_s \sin(2\theta_s)}{f_m}$$

$$A_{32} = \frac{\Delta m_{31}^2}{4E}$$

$$A_s = \pm \frac{G_F N_n}{2\sqrt{2}}$$

$$\sin 2\theta_s = \frac{2\sqrt{d_\mu d_\tau d_s}}{(1 - d_\mu)}$$


$$\cos 2\theta_s = \frac{d_\tau - d_\mu d_s}{(1 - d_\mu)}$$

Our Matter Effect Model

$$P_{\mu\mu} = (1 - d_\mu)^2 |\tilde{S}_{22}|^2 + d_\mu^2$$

$$P_{\mu\tau} = (1 - d_\mu)(1 - d_\tau) + (d_\mu(1 - d_\mu) - d_s(1 + d_\mu)) |\tilde{S}_{22}|^2 - \sqrt{d_\mu d_\tau d_s} \left(\tilde{S}_{23} \tilde{S}_{22}^* + \tilde{S}_{23}^* \tilde{S}_{22} \right)$$

$$P_{\mu s} = (1 - d_\mu)(1 - d_s) + (d_\mu d_s - d_\tau) |\tilde{S}_{22}|^2 + \sqrt{d_\mu d_\tau d_s} \left(\tilde{S}_{23} \tilde{S}_{22}^* + \tilde{S}_{23}^* \tilde{S}_{22} \right)$$



$$|\tilde{S}_{22}|^2 = 1 - \sin^2(2\theta_m) \sin^2(f_m L)$$

$$\left(\tilde{S}_{23} \tilde{S}_{22}^* + \tilde{S}_{23}^* \tilde{S}_{22} \right) = -2 \sin(2\theta_m) \cos(2\theta_m) \sin^2(f_m L)$$

Our Parameters: $|U_{\mu 4}|^2$

- Amount mixing between ν_μ and the sterile mass state ν_4
- Primary effect is extra ν_μ disappearance at **all path lengths**
- Is directly comparable to **SBL measurements of ν_μ disappearance** ($\theta_{\mu\mu}$) and indirectly to the MB/LSND appearance signal ($\theta_{\mu e}$)
- With more sterile neutrinos, becomes a more generic parameter d_μ , but our limit is still applicable:

$$d_\mu = \frac{1 - \sqrt{1 - 4A}}{2} ,$$

$$A = (1 - |U_{\mu 4}|^2 - |U_{\mu 5}|^2 - |U_{\mu 6}|^2)(|U_{\mu 4}|^2 + |U_{\mu 5}|^2 + |U_{\mu 6}|^2) \\ + |U_{\mu 4}|^2 |U_{\mu 5}|^2 + |U_{\mu 4}|^2 |U_{\mu 6}|^2 + |U_{\mu 5}|^2 |U_{\mu 6}|^2 .$$

Our Parameters: $|U_{\tau 4}|^2$

- Amount mixing between ν_τ and the sterile mass state ν_4
- Controls $\nu_\mu \rightarrow \nu_\tau$ vs. $\nu_\mu \rightarrow \nu_s$ fraction
 - Previous SK sterile measurements have implicitly limited this parameter
- This parameter \sim scales the size of sterile-NC matter effects
- Also responsible of NC disappearance over long baselines
- Private to long-baseline and atmospheric measurements
 - But still interesting for understanding atmospheric oscillations